

Hepatitis E Prevalence in North American Blood Donors

IPFA May 16 - 17, Athens

Margaret Fearon, MD, Medical Director, Medical Microbiology

Canadian Blood Services



Disclosures

- Funding for all laboratory support and reagents provided by:
 - Roche Molecular Diagnostics
 - National Microbiology Laboratory (NML), Public Health Agency of Canada (Canadian Study #1)
- Testing performed by:
 - Screening: American Red Cross Scientific Affairs
 - Confirmatory: Sanquin, The Netherlands
 - Sub-population antibody screening: CDC
 - NML (Canadian Study #1)
- Samples provided by
 - American Red Cross
 - Hema-Quebec and Canadian Blood Services



Public Health Agency of
Canada



Sanquin



Collaborators



Canadian Blood Services
Société canadienne du sang

Canadian Blood Services

M. Fearon
S. O'Brien
V. Scalia



HÉMA-QUÉBEC

Héma Québec

G. Delage
F. Bernier
Y. Gregoire



**American
Red Cross**

American Red Cross

S. Stramer
M. Nguyen
S. Smith



Roche Molecular Diagnostics

J. Duncan
N. Huynh
J. Stanley
S. Galel




**Public Health Agency of
Canada, National
Microbiology Laboratory,**
A. Andonov



Sanquin
B Hogema

Hepatitis E Prevalence

Geographical Distribution of Hepatitis E virus Genotypes



Hepatitis E: Current Status Perez-Gracia et al. Reviews in Medical Virology 2013;23(6)

> 3 million cases/year, mainly in developing countries
Seroprevalence 2 – 50% in developed countries

Clinical Presentation

Most cases asymptomatic



- **Acute hepatitis**

- more common in males > 65 yrs. and in pregnant women
- Incubation period 2 weeks – 2 months
- Jaundice, fever, malaise, anorexia
- Viremia is transitory – disappears with symptom onset

- **Chronic hepatitis**

- Chronic HEV infection reported in liver and kidney transplant recipients (SOT), those with non-Hodgkins lymphoma, in HIV/AIDS patients, stem-cell transplant (SCT) patients. (all immunosuppressed patients)
- Almost all cases represent autochthonous HEV gt3
- Persistent viremia with titers 10^5 - 10^7 IU/mL
- Evolution to cirrhosis occurs in up to 50%; liver failure

Risk Profile of Hepatitis E Virus from Pigs or Pork in Canada.

[Wilhelm B](#)¹, [Fazil A](#)², [Rajić A](#)¹, [Houde A](#)³, [McEwen SA](#)¹.

[Send to](#)

[Transbound Emerg Dis.](#) 2017 Dec;64(6):1694-1708.

doi: 10.1111/tbed.12582. Epub 2016 Oct.7



Survey of Canadian retail pork chops and pork livers for detection of hepatitis E virus, norovirus, and rotavirus using real time RT-PCR.

[Wilhelm B](#)¹, [Leblanc D](#)², [Houde A](#)², [Brassard J](#)², [Gagné MJ](#)², [Plante D](#)³, [Bellon-Gagnon P](#)³, [Jones TH](#)⁴, [Muehlhauser V](#)⁴, [Janecko N](#)⁵, [Avery B](#)⁶, [Rajić A](#)⁷, [McEwen SA](#)⁷.

[Int J Food Microbiol.](#) 2014 Aug 18;185:33-40. doi: 10.1016/j.ijfoodmicro.2014.05.006. Epub 2014 May 14



**PIGS, PEOPLE AND
HEPATITIS E
Residue Chemistry and
Predictive Microbiology
Research: Wyndmoor, PA**

Hepatitis E and Transfusion Transmission

- Over 30 cases of reported cases of transfusion transmission.
- Cases reported in Japan, U.K., France, Saudi Arabia.
- To date, no reported cases of TT HEV in North America

North American Donor Prevalence Studies 1

In 2013 **CBS and Hema Quebec** in collaboration with the National Microbiology Lab, carried out a geographically limited HEV prevalence study on approximately 14,000 Canadian blood donors.

- Antibody prevalence of 5.8%
- 0 HEV NAT positive donors (pooled testing 48-100)

Fearon M, O'Brien S. et al Hepatitis E in Canadian Blood Donors. Transfusion 2017;57:1420-142

Also in 2013 the **American Red Cross** surveyed approximately 19,000 U.S. blood donors

- 2 HEV NAT positive donors out of 18,829 tested (ID NAT).
- Antibody prevalence of 7.7% of 4,499 tested donors.

Stramer S, Moritz E. et al. Hepatitis E virus: seroprevalence and frequency of viral RNA detection among US blood donors. Transfusion 2016;56:481-488

North American Donor Prevalence Studies 2

- Two independent RNA prevalence studies – target 100,000 donations total
 - US – between Feb 2015 - Feb 2016
 - Canadian - between October 2016 - June 2017
- **50,000 US (ARC)** donations throughout US but focus in Midwest
 - Unlinked
- **50,000 (30,000 CBS and 20,000 HQ)** donations from every province
 - Linked, tested following product release
- Donations tested for HEV RNA by ID-NAT at the ARC, using the Roche **cobas**® HEV test, and the **cobas**® 6800/8800 System (assay not approved for use in the US or Canada)
 - Hemi-nested PCR from amp/detection plates at Roche
- Additional testing on reactive samples performed at Sanquin and included
 - HEV RNA quantitative PCR (74 bp ORF-3; LOD 10.3 IU/mL)
 - HEV genotyping (ORF-1, ORF-2)
 - HEV antibody testing (Wantai)
- Comparative antibody studies done on a subset of US samples at CDC

Testing System



HEV Analytic Sensitivity	
50% LOD (IU/mL) (95% CI)	3.9 (3.4 - 4.3)
95% LOD (IU/mL) (95% CI)	18.6 (15.9 – 22.6)

The **cobas**® HEV test is not approved in the US or Canada but is available in countries that accept the CE mark.

US Results

3 positives/50,724 valid results
1:16,908 (95% CI: 1:5786-1:81,987)

Sex	Age	State	cobas® HEV Test (Ct value; max = 50)	hn-PCR (Roche)	Viral Load (IU/mL) 2 reps	Geno- type	IgM (S/CO)	IgG (S/CO)
Male	65	IN	38.54	Pos	15, 30	3a	0.02	0.02
Male	21	CA	37.38	Pos	13, 58	3b	0.02	0.02
Female	55	KY	34.42	Pos	1440, 1400	n/a	11.03	>15

Prior published US study (Stramer et al. Transfusion 2016):


2 positives/18,829 valid results

1:9500 (95%CI: 1:2580-1:56,180)

66 yo Male, IA (n/a IU/mL, Ab pos)


71 yo Male, IL (14 IU/mL, Ab neg)

N=3 pos/50,724 (this study)



HEV RNA Positive Donations By Donor Residence

N=2 pos/18,829 (prior study)



Canadian Results

11 positives/50,765 valid results

Total: 1:4615 (95% CI: 1:2579-1:9244)

HQ: 1:2920 (95% CI: 1:1417-1:7262)

CBS: 1:7582 (95% CI: 1:2961-1:27,825)

Follow-up

Site	Sex	Age	Province	cobas® HEV Test (Ct value; max = 50)	Viral load (IU/mL)	IgM (S/CO)	IgG (S/CO)	IgM (S/CO)	IgG (S/CO)
HQ	M	34	Quebec	36.86	68	0.04	0.28	0.05	13.55
HQ	M	60	Quebec	38.92	<10	0.05	1.21	0.1	41.4
HQ	M	59	Quebec	38.89	55	0.01	0.04	0.25	14.62
HQ	M	70	Quebec	38.89	<10	0.47	>15	0.05	276
HQ	F	21	Quebec	34.55	151	2.21	5.87	2.17	31.1
HQ	M	39	Quebec	40.41	28	9.33	14.98	0.91	87.8
HQ	F	41	Quebec	38.29	20	0.01	0.26	2.02	92.4
CBS	M	62	N.S.	37.94	32	0.02	0.03	2.33	147.3
CBS	M	56	Alberta	38.97	43	0.38	0.02	0.76	50.4
CBS	M	17	Alberta	32.35	3080*	0.01	0	0.56	14.91
CBS	M	53	Ontario	39.41	<10	0	0.02	0.1	26.3


***Genotype 3 (unique)**




Subset of 5040 RNA-negative samples tested for IgM/IgG by 3 Commercial HEV antibody assays (CDC; submitted)

Prevalence increases with age

11.4% prevalence
(range 10.65-12.28%)
84% agreement
between assays



1.8% prevalence
(range 0.67-2.9%)
22% agreement
between assays



Summary

- HEV RNA is present in US and Canadian donors
 - Frequency of all ID-NAT studies (Roche/Grifols)
 - 16 pos/120,318 tested, or **1:7500**
 - Wide 95% CIs; no statistical difference amongst studies
 - Highest viral load = 3080 IU/mL; most too low to genotype @ < 100 IU/mL (all gt 3)
 - Most males (13/16)
 - High performance of ID-NAT assays
 - Ab positivity suggests higher frequencies of infection
 - IgM = 1.8% vs IgG = 11.8% => this study
 - IgM = 0.58% vs IgG = 7.7% => prior study
 - IgG = 5.9% Fearon et al; Canada; all significantly different

Summary

- Absence of TT HEV in US/Canada, but clinical disease occurs - magnitude not well understood
 - Kamar et al-chronic HEV after SOT 17-center survey, 2011*; Drobeniuc et al-CDC, surveillance 2005-2012

***“HEV infection causes chronic hepatitis in >60% recipients of solid organ transplants”**

- Hewitt et al-UK; mean viral load
 - **TT = $4.53 \log_{10}$ IU/mL** vs
 - **non-TT = $2.57 \log_{10}$ IU/mL**; 95% CIs overlap
- Decision making underway in Canada
 - RBDM
- Should screened blood be used for:
 - Solid organ, stem cell transplant recipients?

